

HIP-1999-14/TH
KANAZAWA-99-05
March, 1999

Electroweak symmetry breaking and s-spectrum in M-theory

Tatsuo Kobayashi ^{a,b}, Jisuke Kubo ^c and Hitoshi Shimabukuro ^c

^a Department of Physics, High Energy Physics Division
University of Helsinki,

^b Helsinki Institute of Physics
P.O. Box 9, FIN-00014 Helsinki, Finland
and

^c Institute for Theoretical Physics, Kanazawa University
Kanazawa, 920-1192, Japan

Abstract

We study soft SUSY breaking parameters in M-theory with and without 5-brane moduli fields. We investigate successful electroweak symmetry breaking and the positivity of stau mass squared. We study mass spectra in allowed regions.

1 Introduction

Supersymmetry (SUSY) breaking is very important to connect superstring theory to low-energy physics. Induced soft SUSY breaking terms are significant to currently running and future experiment, and they also have cosmological implications. Actually, soft SUSY breaking terms have been derived, e.g. within the framework of weakly coupled heterotic string theory under the assumption that only F -terms of the dilaton S and moduli fields T^i contribute to SUSY breaking [1, 2, 3, 4]. Also various phenomenological aspects of these string-inspired soft terms have been studied.

Recently strongly coupled superstring theories as well as strongly coupled SUSY gauge theories have been studied. Strongly coupled $E_8 \times E_8$ heterotic string theory has been considered as M-theory compactified on S^1/Z_2 [5]. In particular, the 4-dimensional effective theory has been obtained and supersymmetry (SUSY) breaking has been studied [6]-[20]. M-inspired soft SUSY breaking parameters have a certain difference from those derived from weakly coupled string models. Soft parameter formulae between the weakly and strongly coupled regions are connected by the parameter $\alpha(T + \bar{T})$. Actually, several phenomenological aspects of soft SUSY breaking parameters have been investigated, e.g. radiative electroweak symmetry breaking and s-spectrum [21], although the large $\tan\beta$ region has not been studied extensively.

The standard embedding leads to the gauge group $E_6 \times E'_8$, where the top, bottom and tau Yukawa couplings are unified, that is, we have a large value of $\tan\beta$. In this sense, the large $\tan\beta$ scenario is one natural target to study. When E_6 breaks into the standard model (SM) gauge group $G_{SM} = SU(3)_C \times SU(2)_L \times U(1)_Y$, another type of soft scalar terms appear, i.e., D -term contributions.

On the other hand, non-standard embedding including Wilson line breaking is also possible [22] and that can lead to smaller gauge groups, e.g. $SU(3) \times SU(2) \times U(1)$ or $SU(5)$. Thus, in non-standard embedding a small value of $\tan\beta$ can also be realized. Furthermore, non-standard embedding, in general, has 5-branes, that is, in 4-dimensional effective field theory we have a new type of moduli fields Z^n whose vacuum expectation values (VEVs) correspond to positions of 5-branes z^n within $[0, 1]$ along the orbifold S^1/Z_2 . It is possible that F -terms of the 5-brane moduli fields Z^n also contribute to SUSY breaking. Recently 4-dimensional effective theory with Z^n has been

studied [23] and SUSY breaking terms have also investigated [24].

The purpose of this paper is to take into account several effects which are mentioned above and to investigate successful electroweak breaking conditions and the lightest superparticle (LSP) systematically. In the whole paper, we assume that the massless particle content below the grand unified theory (GUT) scale is exactly the same as the minimal supersymmetric standard model (MSSM). In the large $\tan\beta$ scenario, the stau (mass)² becomes small and negative because of large and negative radiative corrections from the large Yukawa coupling. Thus the stau mass leads to a significant constraint for large $\tan\beta$. We cover the whole realistic region of $\tan\beta$ including large $\tan\beta$. We investigate $\alpha(T + \bar{T})$ -dependence of electroweak symmetry breaking and s-spectrum. Furthermore, we study the effect due to F -terms of the moduli fields Z^n .

This paper is organized as follows. In section 2, after we describe the structure of the soft parameters in M-theory without 5-brane, we investigate the electroweak symmetry breaking conditions and the positivity of stau mass squared under the assumption of top-bottom-tau Yukawa coupling unification, that is, the large $\tan\beta$ scenario. D -term contributions, which appear in E_6 breaking into G_{SM} , are also taken into account. We also consider the case with small $\tan\beta$. In section 3, we consider soft parameters in M-theory that the 5-brane moduli field also contribute to SUSY breaking. In this case, electroweak symmetry breaking and mass spectra are studied and allowed regions are compared with the case without 5-brane effects. Section 4 is devoted to conclusions.

2 SUSY breaking in M-theory without 5-branes

In this section we study SUSY breaking in M-theory without 5-branes. For example, standard embedding corresponds to this class. We discuss the soft breaking parameters derived within this framework and study their phenomenological implications.

2.1 Soft parameters

The 4-dimensional effective supergravity action of M-theory can be analyzed by expanding it in powers of the two dimensionless variables: $\varepsilon_1 = \kappa^{\frac{2}{3}}\pi\rho/V^{\frac{2}{3}}$

and $\varepsilon_2 = \kappa^{\frac{2}{3}}/\pi\rho V^{\frac{1}{3}}$, likewise the weakly coupled heterotic string theory expanded by the string coupling $\varepsilon_s = e^{2\phi}/(2\pi)^5$ and the worldsheet sigma model coupling $\varepsilon_\sigma = 4\pi\alpha'/V^{1/3}$, where κ^2 , ρ and V denote the 11-dimensional gravitational coupling, the 11-dimensional length and the Calabi-Yau volume respectively.

Here we consider only the overall moduli field T . Thus, the 4-dimensional effective supergravity includes the dilaton S and the moduli field T in addition to chiral multiplets Φ^p , gauge multiplets and the graviton multiplet. Scalar components of S and T can be identified as

$$\begin{aligned} Re(S) &= \frac{1}{2\pi}(4\pi\kappa^2)^{-2/3}V, \\ Re(T) &= \frac{6^{1/3}}{8\pi}(4\pi\kappa^2)^{-1/3}\pi\rho V^{1/3}. \end{aligned} \quad (1)$$

The Kähler potential is computed by M-theory expansion for matter fields up to order of $\kappa^{4/3}$ [15],

$$K = -\log(S + \bar{S}) - 3\log(T + \bar{T}) + \left(\frac{3}{T + \bar{T}} + \frac{\alpha}{S + \bar{S}}\right)|\Phi|^2. \quad (2)$$

Similarly the gauge kinetic functions of the observable and hidden sectors are calculated [6, 11, 8, 12],

$$f_{E_6} = S + \alpha T, \quad f_{E_8} = S - \alpha T, \quad (3)$$

where $\alpha = 1/(8\pi^2) \int \omega \wedge [tr(F \wedge F) - \frac{1}{2}tr(R \wedge R)]$. In addition, the tree-level superpotential is obtained,

$$W = Y_{pqr}\Phi^p\Phi^q\Phi^r. \quad (4)$$

Here α is of the order one at the “physical” point where gauge coupling unification can be realized [6]. Note that the superpotential W and gauge kinetic functions f_{E_6} and f_{E_8} are exact up to nonperturbative corrections, while there are small additional perturbative corrections to the Kähler potential which are of order $1/4\pi^2 Re(S)$ or $1/[4\pi^2 Re(T)]^3 \sim O(\alpha_{GUT}/\pi)$ in the strong coupling limit $\varepsilon_s \gg 1$.

Now we know the forms of the Kähler potential K , the gauge kinetic function f_{E_6} and the tree-level superpotential W . Thus, we can derive the

formula of soft SUSY breaking parameters following ref.[25, 2]. We assume that a nonperturbative superpotential of S and T is induced and F -terms of S and T contribute to SUSY breaking.

The gravitino mass $m_{3/2}$ is obtained

$$3m_{3/2}^2 = \frac{|F^S|^2}{3(S + \bar{S})^2} + \frac{|F^T|^2}{(T + \bar{T})^2} - V_0, \quad (5)$$

where V_0 is the vacuum energy. Hence, following ref.[3], we parameterize F -terms,

$$F^S = \sqrt{3}m_{3/2}C(S + \bar{S})\sin\theta e^{-i\gamma_S}, \quad (6)$$

$$F^T = m_{3/2}C(T + \bar{T})\cos\theta e^{-i\gamma_T}, \quad (7)$$

where $C^2 = 1 + V_0/3m_{3/2}^2$. Using these parameters, we can write the soft parameters, i.e., the gaugino mass $M_{1/2}$, the soft scalar mass m and the A -parameter as follows [16]

$$M_{1/2} = \frac{\sqrt{3}Cm_{3/2}}{(S + \bar{S}) + \alpha(T + \bar{T})} \left((S + \bar{S})\sin\theta e^{-i\gamma_S} + \frac{\alpha(T + \bar{T})}{\sqrt{3}}\cos\theta e^{-i\gamma_T} \right), \quad (8)$$

$$m^2 = V_0 + m_{3/2}^2 - \frac{3C^2m_{3/2}^2}{3(S + \bar{S}) + \alpha(T + \bar{T})} \times \left\{ \alpha(T + \bar{T}) \left(2 - \frac{\alpha(T + \bar{T})}{3(S + \bar{S}) + \alpha(T + \bar{T})} \right) \sin^2\theta + (S + \bar{S}) \left(2 - \frac{3(S + \bar{S})}{3(S + \bar{S}) + \alpha(T + \bar{T})} \right) \cos^2\theta - \frac{2\sqrt{3}\alpha(T + \bar{T})(S + \bar{S})}{3(S + \bar{S}) + \alpha(T + \bar{T})} \sin\theta \cos\theta \cos(\gamma_S - \gamma_T) \right\}, \quad (9)$$

$$A = \sqrt{3}Cm_{3/2} \left\{ \left(-1 + \frac{3\alpha(T + \bar{T})}{3(S + \bar{S}) + \alpha(T + \bar{T})} \right) \sin\theta e^{-i\gamma_S} + \sqrt{3} \left(-1 + \frac{3(S + \bar{S})}{3(S + \bar{S}) + \alpha(T + \bar{T})} \right) \cos\theta e^{-i\gamma_T} \right\}. \quad (10)$$

In following analyses we concentrate to the case with the vanishing vacuum energy, $V_0 = 0$ ($C = 1$) and no CP phase $\gamma_S = \gamma_T = 0$.

Since the gauge kinetic function f_{E_6} provides the gauge coupling constant, that is, $Ref_{E_6} = 1/g_{E_6}^2$, the phenomenological requirement $g_{GUT}^{-2} \simeq 2$ leads to

$$(S + \bar{S}) + \alpha(T + \bar{T}) \simeq 4. \quad (11)$$

Moreover, the constraint for $Ref_{E_8} > 0$ leads to $(S + \bar{S}) > \alpha(T + \bar{T})$. Therefore, we have

$$0 < \alpha(T + \bar{T}) \lesssim 2. \quad (12)$$

Note that there are three free parameters $m_{3/2}$, θ and $\alpha(T + \bar{T})$, and in particular, $\alpha(T + \bar{T})$ is the characteristic parameter of M-theory. The limit $\alpha(T + \bar{T}) \rightarrow 0$ corresponds to weakly coupled theory.

2.2 Electroweak symmetry breaking and mass spectrum

We now analyze phenomenology of M-theory by using eqs.(8),(9),(10) as boundary conditions of renormalization group equations (RGEs) at the GUT scale. We investigate successful radiative electroweak symmetry breaking and positive masses squared for any sfermion. That gives constraints for the parameters $m_{3/2}$, $\alpha(T + \bar{T})$ and θ .

If we assume a certain type of μ -term generation mechanism, we could fix magnitudes of the supersymmetric Higgs mixing mass μ and the corresponding SUSY breaking parameter B . However, we do not take such a procedure here. Because we would like to study generic case. We determine these magnitudes by using the following minimization conditions of the MSSM Higgs potential,

$$\begin{aligned} m_1^2 + m_2^2 &= -\frac{2\mu B}{\sin \beta}, \\ m_1^2 - m_2^2 &= -(M_Z^2 + m_1^2 + m_2^2) \cos 2\beta, \end{aligned} \quad (13)$$

where $m_{1,2}^2 = m_{H_1, H_2}^2 + \mu^2$, and m_{H_1, H_2}^2 are soft scalar masses of Higgs fields. The Higgs fields for the down sector and the up sector are denoted by H_1 and H_2 . Using these equations, μ and B are written in terms of the soft scalar masses and $\tan \beta$.

Realization of successful electroweak symmetry breaking leads to the condition,

$$m_1^2 m_2^2 < |\mu B|^2, \quad (14)$$

and the bounded-from-below condition along the D -flat directions in the Higgs potential, that is,

$$m_1^2 + m_2^2 > 2|\mu B|, \quad (15)$$

should be satisfied in order for stability of the potential along the D -flat direction. In the large $\tan\beta$ scenario, these conditions lead to

$$m_{H_1}^2 - m_{H_2}^2 \gtrsim M_Z^2/2. \quad (16)$$

Also, we require that a physical mass squared should be positive for any sfermion. Since the sleptons have no large positive-definite radiative corrections from loops involving the gluino, the sleptons become lighter than squarks at low energy. In particular, in the case of large $\tan\beta$, the stau mass squared has a sizable and negative radiative correction due to the large Yukawa coupling. lighter stau mass becomes significant for large $\tan\beta$ *. Furthermore, what is the LSP is an important issue. One candidate is the lightest neutralino and another candidate is the lighter stau for large $\tan\beta$ from the above reason †. The stau mass matrix is

$$\mathbf{m}_{\tilde{\tau}}^2 = \begin{pmatrix} m_{\tilde{\tau}_L}^2 + (-\frac{1}{2} + \sin^2 \theta_W) M_Z^2 \cos 2\beta & v Y_\tau (A_\tau \cos \beta - \mu \sin \beta) \\ v Y_\tau (A_\tau \cos \beta - \mu \sin \beta) & m_{\tilde{\tau}_R}^2 - M_Z^2 \sin^2 \theta_W \cos 2\beta \end{pmatrix}, \quad (17)$$

where $v^2 \equiv \langle H_1 \rangle^2 + \langle H_2 \rangle^2$, and we have neglected the tau (mass)² M_τ^2 term here. The lighter stau mass squared is obtained as

$$\begin{aligned} m_{\tilde{\tau}_1}^2 &= \frac{1}{2} \left(m_{\tilde{\tau}_L}^2 + m_{\tilde{\tau}_R}^2 - \frac{1}{2} M_Z^2 \cos 2\beta \right. \\ &\quad \left. - [(m_{\tilde{\tau}_L}^2 - m_{\tilde{\tau}_R}^2 - (\frac{1}{2} - 2 \sin^2 \theta_W) M_Z^2 \cos 2\beta)^2 \right. \\ &\quad \left. + 4v Y_\tau (A_\tau \cos \beta - \mu \sin \beta)^2]^{1/2} \right). \end{aligned} \quad (18)$$

* For example, in refs.[26, 27] the constraint $m_{\tilde{\tau}_1}^2 > 0$ has been considered in details for (finite) $SU(5)$ GUTs and $SO(10)$ GUTs.

†In ref.[28] cosmological implications of the stau LSP have been discussed. The stau LSP could not be a candidate for cold dark matter. In the case with the stau LSP we need another candidate for cold dark matter.

We take the following input parameters,

$$\begin{aligned}
M_\tau &= 1.777\text{GeV}, & M_Z &= 91.188\text{GeV}, \\
\alpha_{EM}^{-1}(M_Z) &= 127.9 + \frac{8}{9\pi} \log \frac{M_t}{M_Z}, \\
\sin^2 \theta_W &= 0.2319 - 3.03 \times 10^{-5} \Delta_t - 8.4 \times 10^{-8} \Delta_t^2,
\end{aligned} \tag{19}$$

where $\Delta_t = M_t[\text{GeV}] - 165.0$. Here M_τ and M_t are physical tau lepton and top quark masses. We define the SUSY scale as the arithmetic average of the stop squared mass eigenvalues [29],

$$M_{SUSY}^2 = \frac{m_{\tilde{t}_1}^2 + m_{\tilde{t}_2}^2}{2}. \tag{20}$$

2.2.1 Standard embedding

First of all, we consider the case of standard embedding, where the gauge group $E_6 \times E'_8$ is obtained and the top, bottom and tau Yukawa couplings are unified at the GUT scale. From the experimental value of M_τ , we obtain $\tan\beta = 53$ and $M_t = 175\text{GeV}$. We assume at the GUT scale E_6 breaks into G_{SM} and exactly the MSSM matter content is obtained. On top of that, we assume this symmetry breaking induces no extra contributions to soft parameters. Then we use eqs.(8),(9),(10) as the initial conditions of RGEs at the GUT scale. The latter assumption will be relaxed later.

Before studying the constraints, it is useful to show how the soft parameters depend on the parameters, $2\alpha Re(T)$ and θ . The A -parameter is always negative. Fig. 1 shows the ratios $-A/M_{1/2}$ and $m/M_{1/2}$ against $2\alpha Re(T)$ for values of θ . The three solid curves show $-A/M_{1/2}$. The upper, intermediate and lower curves correspond to $-A/M_{1/2}$ for $\theta = 0, \pi/5$ and $\pi/2$, respectively. Similarly, the two dotted curves show $m/M_{1/2}$. The upper and lower correspond to $m/M_{1/2}$ for $\theta = \pi/5$ and $\pi/2$, respectively. The weak coupling limit leads to

$$\frac{A}{M_{1/2}} = -1, \quad \frac{m}{M_{1/2}} = \frac{1}{\sqrt{3}}, \tag{21}$$

for any value of θ [‡]. We have $-A/M_{1/2} \approx -1$ for any value of $2\alpha Re(T)$ except the dilaton-dominant case $|\sin \theta| \approx 1$ and the moduli-dominant case

[‡]These relations have an implication for finiteness and RG invariance [30, 26].

$|\cos \theta| \approx 1$. In the dilaton-dominant (moduli-dominant) case, $-A/M_{1/2}$ becomes smaller (larger) in particular at $2\alpha Re(T) > 1.5$. The ratio $m/M_{1/2} = 1/\sqrt{3}$ holds approximately for any value of $2\alpha Re(T)$ except in the case with $|\sin \theta| > 0.9$ and the purely moduli-dominant case with $|\cot \theta| > O(10)$. In the moduli dominant case with $|\sin \theta| > 0.9$, as $2\alpha Re(T)$ increases, $m/M_{1/2}$ becomes small in particular at $2\alpha Re(T) > 1.5$. Thus, we have similar results in the region where the relations (21) hold approximately.

Figs. 2 show the regions excluded by the electroweak symmetry breaking conditions and the constraint $m_{\tilde{\tau}_1}^2 > 0$. The dotted region denotes the forbidden parameter region by the electroweak breaking conditions, while the squares correspond to $m_{\tilde{\tau}_1}^2 < 0$. In addition the asterisks denote the region with $m_{\tilde{\tau}_1}^2 < m_{\tilde{\chi}_1^0}^2$, where the LSP is the stau.

The constraint due to successful electroweak symmetry breaking is relaxed more as $2\alpha Re(T)$ increases. In particular, a large θ within $[0, \pi]$ is favorable for successful electroweak symmetry breaking. On the other hand, in the case with a large θ , we have $m_{\tilde{\tau}_1}^2 < 0$ for large $2\alpha Re(T)$. That is because we have $m^2 < 0$ at the GUT scale by eq.(9), e.g. in the cases with $\theta = \pi/2$ and $3\pi/4$ for $\alpha(T + \bar{T}) > 1.8$ and 1.3, respectively. In any allowed region, the LSP is the stau.

2.2.2 D -term contributions

We have assumed that symmetry breaking has no effect on soft parameters. However, if a gauge group symmetry breaks reducing its rank, additional contributions to soft scalar mass terms, in general, appear, i.e. D -term contribution [31, 32]. Thus, when the gauge group E_6 breaks to G_{SM} , the additional contributions to soft scalar mass are able to appear at the breaking scale. Here we take into such D -term contributions effects to soft scalar masses. In general, these D -term contributions are proportional to quantum numbers of broken diagonal generators. In the present case, we have two independent universal factors of D -term contributions, which we denote as m_{D1}^2 and m_{D2}^2 , because the rank of E_6 is larger than G_{SM} by two. We regard these as free parameters and then analyze their effects on the parameter space. The soft scalar masses at the GUT scale including D -term contributions are written as

$$m_Q^2 = m_t^2 = m_{\tilde{\tau}_R}^2 = m_{27}^2 - m_{D1}^2 + m_{D2}^2,$$

$$\begin{aligned}
m_b^2 = m_{\tilde{\tau}_L}^2 &= m_{27}^2 + 3m_{D1}^2 + m_{D2}^2, \\
m_{H1}^2 &= m_{27}^2 - 2m_{D1}^2 - 2m_{D2}^2, \\
m_{H2}^2 &= m_{27}^2 + 2m_{D1}^2 - 2m_{D2}^2,
\end{aligned} \tag{22}$$

where m_{27}^2 is the soft scalar mass in eq.(9).

We now analyze D -term contribution effects on the parameter space fixing $\tan \beta = 53$, which has been obtained under the assumption of the top-bottom-tau Yukawa coupling unification. The term m_{D1}^2 with a negative and sizable value is helpful to create a positive gap $m_{H1}^2 - m_{H2}^2$, which is favorable to successful electroweak symmetry breaking. Also, a negative value of m_{D1}^2 increases $m_{\tilde{\tau}_R}^2$, but it reduces $m_{\tilde{\tau}_L}^2$. The (mass)² $m_{\tilde{\tau}_R}^2$ is dominant to $m_{\tilde{\tau}_1}^2$ around the universal case with $m_{\tilde{\tau}_R}^2 = m_{\tilde{\tau}_L}^2$. Therefore negative m_{D1}^2 may increase $m_{\tilde{\tau}_1}^2$ and seems to be favorable to the constraint of $m_{\tilde{\tau}_1}^2$. However, as m_{D1}^2 decreases, $m_{\tilde{\tau}_L}^2$ is decreased and below a critical value of m_{D1}^2 $m_{\tilde{\tau}_L}^2$ becomes dominant to $m_{\tilde{\tau}_1}^2$. A similar effect of the D -term contribution has been discussed within the framework of $SO(10)$ GUTs [27].

On the other hand, a positive value of m_{D2}^2 increases both $m_{\tilde{\tau}_R}^2$ and $m_{\tilde{\tau}_L}^2$. Thus, a positive value of m_{D2}^2 is favorable to increase $m_{\tilde{\tau}_1}^2$. The effects of m_{D2}^2 to m_{H1}^2 and m_{H2}^2 are same. Hence, m_{D2}^2 has little effect on electroweak symmetry breaking.

These results are shown in Figs.3 and Figs. 4 for $\theta = \pi/2$. Figs. 3 show the m_{D2}^2 effects. A large value of m_{D2}^2 reduces the region excluded by the constraint $m_{\tilde{\tau}_1}^2 > 0$, while it has little effect on electroweak symmetry breaking. The region with the neutralino LSP appears and it becomes wider as m_{D2}^2 increases. For example, in the case of $m_{D2}^2 = 1.0 \text{ (TeV)}^2$, the region with the neutralino LSP has the largest value $m_{3/2} = 3.1 \text{ TeV}$ when $\alpha(T + \bar{T}) = 0.8$. In this case the lightest neutral Higgs boson mass m_{h^0} is 151 GeV. Here we use the two loop Higgs mass m_{h^0} [29, 33]. Similarly, we obtain $m_{h^0} = 129 \text{ GeV}$ when $m_{3/2} = 1.0 \text{ TeV}$ and $\alpha(T + \bar{T}) = 0.8$. This value is stable between $m_{3/2} = 0.4 \text{ TeV}$ and 1.0 TeV .

Figs. 4 show the m_{D1}^2 effects. They show that the region excluded by the electroweak symmetry breaking conditions becomes more narrow and the region with $m_{\tilde{\tau}_1}^2 < 0$ becomes wider as m_{D1}^2 decreases. In suitable values of m_{D1}^2 , we have the region with the neutralino LSP. Figs. 4 show the critical value is around $m_{D1}^2 = -2.0 \text{ (TeV)}^2$, where the neutralino LSP region is widest.

2.2.3 Small $\tan\beta$ scenario

It is also possible that we obtain a smaller gauge group, e.g. G_{SM} or $SU(5)$ at the GUT scale by non-standard embedding including Wilson line breaking. In this case, the assumption of the top-bottom-tau Yukawa coupling unification can be relaxed and a smaller value of $\tan\beta$ can be realized. We assume that below the GUT scale the matter content is exactly same as the MSSM. Here, we vary $\tan\beta$ holding eqs.(8),(9),(10) as the initial conditions of RGEs at the GUT scale and fixing $M_t = 175$ GeV.

Figs. 5 show that the allowed parameter region for $\tan\beta$. Fig. 5a shows that for a small value of $|\sin\theta|$ we have no excluded region at $\tan\beta < 50$. However, for large values of $|\sin\theta|$ and $2\alpha Re(T)$, both the regions with $m_{\tilde{\tau}_1}^2 < 0$ and the stau LSP remain even when $\tan\beta$ is small as shown in Fig. 5b.

3 M-theory with 5-brane

3.1 soft parameters

In general, non-standard embedding includes 5-branes, that is, 4-dimensional effective theory includes 5-brane moduli fields Z^n . It is possible that Z^n also contribute to SUSY breaking. Thus, we consider such effects for generic case. If the 5-branes exist between the boundaries in M-theory, the Kähler potential and the gauge kinetic functions are modified. The relevant part of the moduli Kähler potential K_{mod} and the gauge kinetic functions of the observable and hidden sectors $f^{(1)}$ and $f^{(2)}$ with N 5-branes take the following form [23, 24],

$$K_{mod} = -\ln(S + \bar{S}) - 3\ln(T + \bar{T}) + K_5, \quad (23)$$

$$\begin{aligned} f^{(1)} &= S + \epsilon T \left(\beta^{(0)} + \sum_{n=1}^N (1 - Z^n)^2 \beta^{(n)} \right), \\ f^{(2)} &= S + \epsilon T \left(\beta^{(N+1)} + \sum_{n=1}^N (Z^n)^2 \beta^{(n)} \right), \end{aligned} \quad (24)$$

where K_5 is the Kähler potential for the 5-brane position moduli Z^n and the expansion parameter $\epsilon = (\kappa/4\pi)^{2/3} 2\pi^2 \rho/V^{2/3}$. In addition, $\beta^{(0)}$, $\beta^{(N+1)}$ are the instanton numbers on the observable sector boundary and the hidden

sector boundary respectively, and $\beta^{(n)}$ ($n = 1, \dots, N$) is a magnetic charge on the each 5-brane. They should satisfy the cohomology constraint,

$$\sum_{n=0}^{N+1} \beta_i^{(n)} = 0, \quad (25)$$

which means physically that the net charges must be zero because the “flux” cannot get away anywhere in compact space.

The part of the matter fields is obtained

$$K_{mat} = \left(\frac{3}{(T + \bar{T})} + \frac{\epsilon\zeta}{(S + \bar{S})} \right) |\Phi|^2, \quad (26)$$

where

$$\zeta = \beta^{(0)} + \sum_{n=1}^N (1 - z^n)^2 \beta^{(n)}. \quad (27)$$

Following the generic formulae of refs.[25, 2] again, we can derive the soft parameters. We assume $V_0 = 0$ again. The soft parameters are obtained,

$$\begin{aligned} M_{1/2} &= \frac{1}{(S + \bar{S}) + \epsilon\zeta(T + \bar{T})} [F^S + \epsilon\zeta F^T + \epsilon T F^n \zeta_n] \\ m^2 &= m_{3/2}^2 - |F^S|^2 \left[\frac{1}{(S + \bar{S})^2} - \frac{9}{(3(S + \bar{S}) + \epsilon\zeta(T + \bar{T}))^2} \right] \\ &\quad - |F^T|^2 \left[\frac{1}{(T + \bar{T})^2} - \frac{(\epsilon\zeta)^2}{(3(S + \bar{S}) + \epsilon\zeta(T + \bar{T}))^2} \right] \\ &\quad - F^n \bar{F}^{\bar{m}} \left[\frac{\epsilon\zeta_{n\bar{m}}(T + \bar{T})}{3(S + \bar{S}) + \epsilon\zeta(T + \bar{T})} - \frac{\epsilon^2 \zeta_n \zeta_{\bar{m}} (T + \bar{T})^2}{(3(S + \bar{S}) + \epsilon\zeta(T + \bar{T}))^2} \right] \\ &\quad + 3F^S \bar{F}^{\bar{T}} \frac{\epsilon\zeta}{(3(S + \bar{S}) + \epsilon\zeta(T + \bar{T}))^2} + 3F^S \bar{F}^{\bar{m}} \frac{\epsilon\zeta_{\bar{m}}(T + \bar{T})}{(3(S + \bar{S}) + \epsilon\zeta(T + \bar{T}))^2} \\ &\quad - F^T \bar{F}^{\bar{m}} \left[\frac{\epsilon\zeta_{\bar{m}}}{3(S + \bar{S}) + \epsilon\zeta(T + \bar{T})} - \frac{\epsilon^2 \zeta \zeta_{\bar{m}} (T + \bar{T})}{(3(S + \bar{S}) + \epsilon\zeta(T + \bar{T}))^2} \right] + (c.c) \end{aligned} \quad (28)$$

$$(29)$$

$$\begin{aligned}
A &= F^S \left[\frac{2}{S + \bar{S}} - \frac{9}{3(S + \bar{S}) + \epsilon \zeta (T + \bar{T})} \right] \\
&+ F^T \frac{\epsilon \zeta}{3(S + \bar{S}) + \epsilon \zeta (T + \bar{T})} \\
&+ F^n \left[K_{5,n} - \frac{3\epsilon \zeta_n (T + \bar{T})}{3(S + \bar{S}) + \epsilon \zeta (T + \bar{T})} \right],
\end{aligned} \tag{30}$$

where $K_{5,n} = \partial K_5 / \partial Z^n$, $\zeta_n = \partial \zeta / \partial z^n$ and $\zeta_{nm} = \partial^2 \zeta / (\partial z^n \partial z^m)$.

We have normalized the soft scalar mass as,

$$m_{IJ}^2 = m^2 \frac{\partial^2 K_{mat}}{|\partial \Phi|^2}, \tag{31}$$

using the total Kähler metric. That is different from ref.[24], where the soft scalar mass is normalized as,

$$m_{IJ}^2 = m^2 \frac{3}{(T + \bar{T})^2}. \tag{32}$$

In addition, here we have kept all terms, although in ref.[24] only the linear terms of ε have been kept.

Here we ignore CP phases, i.e. $F = \bar{F}$. For simplicity, we assume that there is only one relevant 5-brane moduli Z and its Kähler potential is a function of only $(Z + \bar{Z})$, i.e.,

$$K_5 = K_5(Z + \bar{Z}). \tag{33}$$

The gravitino mass is obtained,

$$m_{3/2}^2 = \frac{|F^S|^2}{3(S + \bar{S})^2} + \frac{|F^T|^2}{(T + \bar{T})^2} + \frac{1}{3} K_{5,Z\bar{Z}} |F^Z|^2. \tag{34}$$

Now we can parameterize each F -term as follows

$$\begin{aligned}
F^S &= \sqrt{3} m_{3/2} (S + \bar{S}) \sin \theta \sin \phi, \\
F^T &= m_{3/2} (T + \bar{T}) \cos \theta \sin \phi, \\
F^Z &= \sqrt{\frac{3}{K_{5,Z\bar{Z}}}} m_{3/2} \cos \phi.
\end{aligned} \tag{35}$$

Still now we have several unknown parameters and most of them are due to detailed information of the 5-brane. Thus, we fix z and $\beta^{(1)}$, e.g.

$$z = \frac{1}{2}, \quad \beta^{(1)} = \frac{4}{3}\beta^{(0)}. \quad (36)$$

Note that both the instanton number $\beta^{(0)}$ and the intersection number $\beta^{(1)}$ are integers. Eq.(36) leads to the following simple relations with eq.(27) as

$$\zeta|_{z=1/2} = -\zeta_{,z}|_{z=1/2} = \frac{1}{2}\zeta_{,zz}|_{z=1/2} = \beta^{(1)}. \quad (37)$$

In this case we can rewrite the soft parameters,

$$\begin{aligned} M_{1/2} &= \frac{m_{3/2}}{(S + \bar{S}) + \epsilon\zeta(T + \bar{T})} \left\{ \sqrt{3}(S + \bar{S}) \sin \theta \sin \phi \right. \\ &\quad \left. + \epsilon\zeta(T + \bar{T}) \cos \theta \sin \phi - \frac{\epsilon\zeta(T + \bar{T})}{2} \sqrt{\frac{3}{K_{5,Z\bar{Z}}}} \cos \phi \right\}, \\ m^2 &= m_{3/2}^2 \left\{ 1 - 3 \left[1 - \frac{9(S + \bar{S})^2}{(3(S + \bar{S}) + \epsilon\zeta(T + \bar{T}))^2} \right] \sin^2 \theta \sin^2 \phi \right. \\ &\quad - \left[1 - \left(\frac{\epsilon\zeta(T + \bar{T})}{(3(S + \bar{S}) + \epsilon\zeta(T + \bar{T}))} \right)^2 \right] \cos^2 \theta \sin^2 \phi \\ &\quad - \frac{3\epsilon\zeta(T + \bar{T})}{K_{5,Z\bar{Z}}(3(S + \bar{S}) + \epsilon\zeta(T + \bar{T}))} \left[2 - \frac{\epsilon\zeta(T + \bar{T})}{(3(S + \bar{S}) + \epsilon\zeta(T + \bar{T}))} \right] \cos^2 \phi \\ &\quad + \frac{6\sqrt{3}(S + \bar{S})\epsilon\zeta(T + \bar{T})}{(3(S + \bar{S}) + \epsilon\zeta(T + \bar{T}))^2} \sin \theta \cos \theta \sin \phi \\ &\quad - \frac{6\sqrt{3}(S + \bar{S})\epsilon\zeta(T + \bar{T})}{(3(S + \bar{S}) + \epsilon\zeta(T + \bar{T}))^2} \sqrt{\frac{3}{K_{5,Z\bar{Z}}}} \sin \theta \sin \phi \cos \phi \\ &\quad + \frac{2\epsilon\zeta(T + \bar{T})}{3(S + \bar{S}) + \epsilon\zeta(T + \bar{T})} \left[1 - \frac{\epsilon\zeta(T + \bar{T})}{(3(S + \bar{S}) + \epsilon\zeta(T + \bar{T}))} \right] \\ &\quad \left. \times \sqrt{\frac{3}{K_{5,Z\bar{Z}}}} \cos \theta \sin \phi \cos \phi \right\}, \end{aligned}$$

$$\begin{aligned}
A = & m_{3/2} \left\{ \sqrt{3} \left[2 - \frac{9(S + \bar{S})}{3(S + \bar{S}) + \epsilon\zeta(T + \bar{T})} \right] \sin \theta \sin \phi \right. \\
& - \frac{3\epsilon\zeta(T + \bar{T})}{3(S + \bar{S}) + \epsilon\zeta(T + \bar{T})} \cos \theta \sin \phi \\
& + \left. \left[K_{5,Z} + \frac{3\epsilon\zeta(T + \bar{T})}{3(S + \bar{S}) + \epsilon\zeta(T + \bar{T})} \right] \sqrt{\frac{3}{K_{5,Z\bar{Z}}}} \cos \phi \right\}, \tag{38}
\end{aligned}$$

where we have assumed $T = (T + \bar{T})/2$.

In addition, from the gauge kinetic functions (24) we obtain constraints for values of moduli S and T as,

$$2Re(f^{(1)}) = (S + \bar{S}) + \epsilon\zeta(T + \bar{T}) = 2g_{GUT}^{-2} \simeq 4, \tag{39}$$

and

$$2Re(f^{(2)}) = (S + \bar{S}) - \frac{3}{2}\epsilon\zeta(T + \bar{T}) > 0. \tag{40}$$

Here we use eqs.(36), (37), (25), (39) and (40), so that we find the parameter region as $0 < \epsilon\zeta(T + \bar{T}) < 8/5$. That is almost comparable to the region $0 < \alpha(T + \bar{T}) < 2$ in the case without 5-branes.

3.2 Electroweak symmetry breaking and mass spectrum

Using eq.(38), we analyze the 5-brane effects on electroweak symmetry breaking and the mass spectrum. Eq.(38) has total 6 free parameters, that is, $m_{3/2}$, $\epsilon\zeta(T + \bar{T})$, θ , ϕ , and $K_{5,Z\bar{Z}}$, $K_{5,Z}$. Here we use the same notation for $\epsilon\zeta(T + \bar{T})$ as the case without 5-brane, i.e., $\alpha(T + \bar{T}) \equiv \epsilon\zeta(T + \bar{T})$. Note that $K_{5,Z\bar{Z}}$ is positive definite value due to eq.(34), (33), but $K_{5,Z}$ is any real number. The parameter $K_{5,Z}$ appears only in the A -parameter and $K_{5,Z}$ -dependence of A is easy to understand, that is, $A/\cos \phi$ increases linearly as $K_{5,Z}$ increases.

Now, let us concentrate ourselves to the region with $|F^S| \gg |F^T|$, that is, $\sin \theta \approx 1$. In this region, eq.(38) is reduced to

$$M_{1/2} = \frac{m_{\frac{3}{2}}}{4} \left\{ \sqrt{3}(4 - \alpha(T + \bar{T})) \sin \phi - \frac{\alpha(T + \bar{T})}{2} \sqrt{\frac{3}{K_{5,Z\bar{Z}}}} \cos \phi \right\},$$

$$\begin{aligned}
m^2 &= m_{3/2}^2 \left\{ 1 - 3 \left[1 - \frac{9(4 - \alpha(T + \bar{T}))^2}{(3(S + \bar{S}) + \alpha(T + \bar{T}))^2} \right] \sin^2 \phi \right. \\
&\quad - \frac{3\alpha(T + \bar{T})}{K_{5,Z\bar{Z}}(3(S + \bar{S}) + \alpha(T + \bar{T}))} \left[2 - \frac{\alpha(T + \bar{T})}{(3(S + \bar{S}) + \alpha(T + \bar{T}))} \right] \cos^2 \phi \\
&\quad \left. - \frac{6\sqrt{3}(4 - \alpha(T + \bar{T}))\alpha(T + \bar{T})}{(3(S + \bar{S}) + \alpha(T + \bar{T}))^2} \sqrt{\frac{3}{K_{5,Z\bar{Z}}}} \sin \phi \cos \phi \right\}, \\
A &= m_{3/2} \left\{ \sqrt{3} \left[2 - \frac{9(4 - \alpha(T + \bar{T}))}{3(S + \bar{S}) + \alpha(T + \bar{T})} \right] \sin \phi \right. \\
&\quad \left. + \left[K_{5,Z} + \frac{3\alpha(T + \bar{T})}{3(S + \bar{S}) + \alpha(T + \bar{T})} \right] \sqrt{\frac{3}{K_{5,Z\bar{Z}}}} \cos \phi \right\}. \tag{41}
\end{aligned}$$

To compare with the results in section 2 without 5-brane, here we take $\tan \beta = 53$. Actually, in the case with such large $\tan \beta$ we have stronger constraints. At first, we consider the case with $K_{5,Z\bar{Z}} = K_{5,Z} = 1$. Fig. 6 shows $M_{1/2}$, m and A for $\phi = \pi/4$ as well as $m_{3/2} = 2.0$ TeV, $\theta = \pi/2$, and $K_{5,Z\bar{Z}} = K_{5,Z} = 1$. Obviously, the ratios, $A/M_{1/2}$ and $m/M_{1/2}$, are different from eq.(21). The A -parameter is always positive when $\phi = \pi/4$. For small $\alpha(T + \bar{T})$, the dominant terms are the first term of $M_{1/2}$, and the second term of m^2 . The 5-brane SUSY breaking effect F^Z within $0 < \sin \phi < 1$ reduces $M_{1/2}$, but increases m^2 . That is obvious from the comparison between Fig. 6 and eq.(21). Therefore we obtain $m_{\tilde{\tau}_1}^2 > m_{\tilde{\chi}_1^0}^2 \sim M_1^2$ at small $\alpha(T + \bar{T})$ and the lightest neutralino is almost Bino-like. When $\alpha(T + \bar{T})$ become larger, the third and forth term of m^2 become more sizable and m^2 itself is decreased. Thus the excluded region by constraint $m_{\tilde{\tau}_1}^2 > 0$ is larger than the case without the 5-brane effect. Such behavior on electroweak symmetry breaking and the constraint $m_{\tilde{\tau}_1}^2 > 0$ is shown in Fig. 7a for $\theta = \pi/2$, $\phi = \pi/4$ and $K_{5,Z\bar{Z}} = K_{5,Z} = 1$. In the allowed region, the lightest neutralino is almost Bino-like. Fig. 7b shows the case with $\phi = \pi/5$. In the 5-brane dominant case, $\cos \phi \rightarrow 1$, most of the region is excluded. Figs. 8a and 8b show $\alpha(T + \bar{T})$ -dependence and ϕ -dependence of representative superparticle masses, e.g. the lightest neutralino, chargino, stop and stau. The ϕ -dependence is rather larger than $\alpha(T + \bar{T})$ -dependence. When we vary ϕ , the lightest Higgs mass change $132 \text{ GeV} < m_{h^0} < 138 \text{ GeV}$ in the allowed

region for $m_{3/2} = 1$ TeV and $\theta = \pi/2$. The lightest Higgs mass is rather stable for change of $\alpha(T + \bar{T})$.

Next, we calculate $K_{5,Z\bar{Z}}$ contribution fixing $K_{5,Z} = 1$. Fig. 9 shows $M_{1/2}$, m and $|A|$ for $m_{3/2} = 2.0$ TeV, $\theta = \pi/2$, $\phi = \pi/4$, $2\alpha Re(T) = 1.0$ and $K_{5,Z} = 1$. Obviously, the $A/M_{1/2}$ and $m/M_{1/2}$, are different from eq.(21), again. The A -parameter changes from positive to negative around $K_{5,Z\bar{Z}} = 2.5$ for $\phi = \pi/4$ and $2\alpha Re(T) = 1.0$. The terms of the 5-brane contribution of $M_{1/2}$ and m^2 are proportional to $\sqrt{3/K_{5,Z\bar{Z}}}$ or $3/K_{5,Z\bar{Z}}$. When $K_{5,Z\bar{Z}}$ increases, then these terms are suppressed and these soft parameters become close to those without 5-brane except for the existence of the “reducing” factor $\sin \phi$. Therefore the effects of $\sin \phi$ is favorable for the $m_{\tilde{\tau}_1}^2 > 0$ condition and it leads to $m_{\tilde{\tau}_1}^2 > m_{\tilde{\chi}_1^0}^2$. Fig. 10 shows such $K_{5,Z\bar{Z}}$ -dependence of the regions excluded by the electroweak symmetry breaking and the constraint $m_{\tilde{\tau}_1}^2 > 0$ as well as the stau LSP region. Actually, large $K_{5,Z\bar{Z}}$ is useful to increase $m_{\tilde{\tau}_1}^2$, but it is not favorable for successful electroweak symmetry breaking.

Finally, we show $K_{5,Z}$ -dependence, which has the linear effect only on A . Figs. 11 show the $K_{5,Z}$ effect on the electroweak symmetry breaking, the constraint $m_{\tilde{\tau}_1}^2 > 0$ and the LSP for $m_{3/2} = 2.0$ TeV, $\phi = \pi/4$ and $K_{5,Z\bar{Z}} = 1$. As $K_{5,Z}$ increases, the electroweak symmetry breaking condition is relaxed, but the constraint $m_{\tilde{\tau}_1}^2 > 0$ becomes severe. Actually, the region with unsuccessful electroweak symmetry breaking disappears at $K_{5,Z} > 1.4$, but at $K_{5,Z} > 3.2$ all the region is excluded because of $m_{\tilde{\tau}_1}^2 < 0$. Fig. 12 shows $K_{5,Z}$ -dependence of representative superparticle masses. The lightest Higgs mass is rather stable against change of $K_{5,Z}$.

4 Conclusions

We have studied soft SUSY breaking parameters in M-theory with and without 5-brane moduli effects. We investigate successful electroweak symmetry breaking and the constraint $m_{\tilde{\tau}_1}^2 > 0$. In the allowed regions, we have shown mass spectra.

We have considered the large $\tan \beta$ scenario, because these constraints are severe for large $\tan \beta$. In the case without 5-brane effects, the electroweak symmetry breaking conditions are relaxed in the strong coupling region $\alpha(T + \bar{T}) > 1.5$. In such region, however, the constraint $m_{\tilde{\tau}_1}^2 > 0$ becomes strong in particular for large θ . In any allowed region, the LSP is the stau.

The positive D -term contribution m_{D2}^2 can relax the constraint $m_{\tilde{\tau}_1}^2 > 0$ and lead to the region with the neutralino LSP. The negative D -term contribution m_{D1}^2 can relax the electroweak symmetry breaking conditions. In the suitable value of m_{D1}^2 we have the widest region with the neutralino LSP.

For small $\tan\beta$, in particular $\tan\beta < 50$, the electroweak symmetry breaking is realized easily. However, for large $\alpha(T + \bar{T})$ and large θ , the region with $m_{\tilde{\tau}_1}^2 < 0$ and the region with the stau LSP remain.

The 5-brane effect can relax the electroweak symmetry breaking condition, but it makes the $m_{\tilde{\tau}_1}^2 > 0$ constraint severe.

Soft terms in M-theory with 5-branes include more free parameters. Thus, we will leave to future work systematic study of the whole parameter space on other phenomenological aspects, e.g. cosmological implications and the $b \rightarrow s\gamma$ decay [34], which leads to another constraint for large $\tan\beta$ and lighter superparticles in the universal case.

Acknowledgement

The authors would like to thank K. Enqvist and S.-J. Rey for useful discussions. This work was partially supported by the Academy of Finland (no. 44129).

References

- [1] L.E. Ibáñez and D. Lüst, Nucl. Phys. **B382** (1992) 305;
B. deCarlos, J.A. Casas and C. Muñoz, Phys. Lett. **B299** (1993) 234.
- [2] V.S. Kaplunovsky and J. Louis, Phys. Lett. **B306** (1993) 269.
- [3] A. Brignole, L.E. Ibáñez and C. Muñoz, Nucl. Phys. **B422** (1994) 125.
- [4] T. Kobayashi, D. Suematsu, K. Yamada and Y. Yamagishi, Phys. Lett. **B348** (1995) 402.
- [5] P. Hořava and E. Witten, Nucl. Phys. **B460** (1996) 506;
Nucl.Phys.**B475**, (1996) 94;

- P. Hořava, Phys. Rev. **D54** (1996) 7561;
E. Witten, Nucl. Phys. **B471** (1996) 135.
- [6] T. Banks and M. Dine, Nucl. Phys. **B479** (1996) 173.
 - [7] I. Antoniadis and M. Quirós, Phys. Lett. **B392** (1997) 61; Phys. Lett. **B416** (1998) 327.
 - [8] H.P. Nilles and S. Stieberger, Nucl. Phys. **B499** (1997) 3.
 - [9] T. Li, J.L. Lopez and D.V. Nanopoulos, Phys. Rev. **D56** (1997) 2602.
 - [10] E. Dudas and C. Grojean, Nucl. Phys. **B507** (1997) 553.
 - [11] K. Choi, Phys. Rev. **D56** (1997) 6588.
 - [12] H.P. Nilles, M. Olechowski and M. Yamaguchi, Phys. Lett. **B415** (1997) 24; Nucl. Phys. **B530** (1998) 43.
 - [13] Z. Lalak and S. Thomas, Nucl. Phys. **B515** (1998) 55.
 - [14] E. Dudas, Phys. Lett. **B416** (1998) 309.
 - [15] A. Lukas, B.A. Ovrut and D. Waldram, Nucl. Phys. **B532** (1998) 43; Phys. Rev. **D57** (1998) 7529.
 - [16] K. Choi, H.B. Kim and C. Muñoz, Phys. Rev. **D57** (1998) 7521.
 - [17] T. Li, Phys. Rev. **D57** (1998) 7539.
 - [18] A. Lukas, B.A. Ovrut, K.S. Stelle and D. Waldram, Phys. Rev. **D59** (1998) 086001; hep-th/9806051.
 - [19] J. Ellis, Z. Lalak, S. Pokorski and W. Pokorski, Nucl. Phys. **B540** (1998) 149.
 - [20] J. Ellis, Z. Lalak and W. Pokorski, hep-th/9811133.
 - [21] D. Bailin, G.V. Kraniotis and A. Love, Phys. Lett. **B432** (1998) 90; hep-ph/9812283;
Y. Kawamura, H.P. Nilles, M. Olechowski and M. Yamaguchi, J. High Energy Phys. **06** (1998) 008;

- S.A. Abel and C.A. Savoy, hep-ph/9809498;
J.A. Casas, A. Ibarra and C. Muñoz, hep-ph/9810266;
C.S. Huang, T. Li, W. Liao and Q.S. Yan and S.H. Zhu, hep-ph/9810412;
T. Li, hep-ph/9903371.
- [22] S. Stieberger, Nucl. Phys. **B541** (1999) 109;
P. Binétury, C. Deffayet, E. Dudas and P. Ramond, Phys. Lett. **B441**
(1998) 163;
Z. Lalak, S. Pokorski and S. Thomas, hep-ph/9807503.
- [23] A. Lukas, B.A. Ovrut and D. Waldram, hep-th/9808101;
R. Donagi, A. Lukas, B.A. Ovrut and D. Waldram, hep-th/9811168;
hep-th/9901009.
- [24] A. Lukas, B.A. Ovrut and D. Waldram, hep-th/9901017.
- [25] S.K. Soni and H.A. Weldon, Phys. Lett. **B126** (1983) 215.
- [26] T. Kobayashi, J. Kubo, M. Mondragón and G. Zoupanos, Nucl. Phys.
B511 (1998) 45.
- [27] Y. Kawamura, T. Kobayashi and H. Shimabukuro, Phys. Lett. **B436**
(1998) 108.
- [28] L.E. Ibáñez, Phys. Lett. **B137** (1984) 160;
J.S. Hagelin, G.L. Kane and S. Raby, Nucl. Phys. **B241** (1984) 638;
T. Falk and K.A. Olive and M. Srednichi, Phys. Lett. **B339** (1994) 248;
L.J. Hall, T. Moroi and H. Murayama, Phys. Lett. **B424** (1998) 305.
- [29] M. Carena, J.R. Espinosa, M. Quirós, C.E.M. Wagner, Phys. Lett.
B355 (1995) 209;
M. Carena, M. Quirós, C.E.M. Wagner, Nucl. Phys. **B461** (1996) 407.
- [30] D.R.T. Jones, L. Mezincescu and Y.P. Yau, Phys. Lett. **B148** (1984)
317;
I. Jack and D.R.T. Jones, Phys. Lett. **B349** (1995) 294;
D.I. Kazakov, M.Yu. Kalmykov, I.N. Kondrashuk and A.V. Gladyshev,
Nucl. Phys. **B471** (1996) 387;
Y. Kawamura, T. Kobayashi and J. Kubo, Phys. Lett. **B405** (1997) 64;
D.I. Kazakov, Phys. Lett. **B421** (1998) 211;

- I. Jack, D.R.T. Jones and A. Pickering, Phys. Lett. **B426** (1998) 73;
T. Kobayashi, J. Kubo and G. Zoupanos, Phys. Lett. **B427** (1998) 291.
- [31] M. Drees, Phys. Lett. **B181** (1986) 279;
J.S. Hagelin and S. Kelley, Nucl. Phys. **B342** (1990) 95;
A.E. Faraggi, J.S. Hagelin, S. Kelley and D.V. Nanopoulos, Phys. Rev. **D45** (1992) 3272;
Y. Kawamura, H. Murayama and M. Yamaguchi, Phys. Lett. **B324** (1994) 54; Phys. Rev. **D51** (1995) 1337;
Y. Kawamura and M. Tanaka, Prog. Theor. Phys. **91** (1994) 949.
- [32] Y. Kawamura, Phys. Rev. **D53** (1996) 3779;
Y. Kawamura and T. Kobayashi, Phys. Lett. **B375** (1996) 141; Phys. Rev. **D56** (1997) 3844;
Y. Kawamura, T. Kobayashi and T. Komatsu, Phys. Lett. **B400** (1997) 284.
- [33] H.E. Haber, R. Hempfling and A.H. Hoang, Z. Phys. **C75** (1997) 539;
S. Heinemeyer, W. Hollik, G. Weiglein, Phys. Rev. **D58** (1998) 091701.
- [34] S. Bertolini, F. Borzumati, A. Masiero and G. Ridolfi, Nucl. Phys. **B353** (1991) 591;
R. Barbieri and G.F. Giudice, Phys. Lett. **B309** (1993) 86;
M. Diaz, Phys. Lett. **B322** (1994) 207;
J.L. López, D.V. Nanopoulos, G.T. Park and A. Zichichi, Phys. Rev. **D49** (1994) 355;
R. Arnowitt and P. Nath, Phys. Lett. **B336** (1994) 395;
B. de Carlos and J.A. Casas, Phys. Lett. **B349** (1995) 300;
S. Bertolini and F. Vissani, Z. Phys. **C67** (1995) 513.

Figure Captions

- Fig. 1 The ratios $-A/M_{1/2}$ and $m/M_{1/2}$ against $2\alpha Re(T)$. The three solid curves correspond to $-A/M_{1/2}$ for $\theta = 0, \pi/5$ and $\pi/2$. The two dotted curves correspond to $m/M_{1/2}$ for $\theta = \pi/5$ and $\pi/2$.
- Fig. 2 The excluded regions by the electroweak breaking condition and the constraint $m_{\tilde{\tau}_1}^2 > 0$. Also the regions with $m_{\tilde{\tau}_1}^2 > m_{\tilde{\chi}_1}^2$ are shown. Fig.2a, 2b, 2c and 2d correspond to the cases with $\theta = \pi/5, \pi/4, \pi/2, 3\pi/4$.
- Fig. 3 Effects of the D -term contribution m_{D2}^2 on the excluded region for $\theta = \pi/2$. Figs.3a and 3b correspond to the cases with $(m_{D1}^2, m_{D2}^2) = (0, 0.5)$ and $(0, 1.0)$ $[(\text{TeV})^2]$, respectively.
- Fig. 4 Effects of the D -term contribution m_{D1}^2 on the excluded region for $\theta = \pi/2$. Figs. 4a, 4b and 4c correspond to the cases with $(m_{D1}^2, m_{D2}^2) = (-0.05, 0), (-2.0, 0)$ and $(-5.0, 0)$ $[(\text{TeV})^2]$, respectively.
- Fig. 5 The excluded region for $\tan\beta$. Fig. 5a corresponds to the case with $\theta = \pi/5$ and $m_{3/2} = 1.0$ TeV for $M_{top} = 175.0$ GeV. Similarly, Fig. 5b corresponds to the case with $\theta = \pi/2$ and $m_{3/2} = 2.0$ TeV for $M_{top} = 175.0$ GeV.
- Fig. 6 The soft parameters, $M_{1/2}$, $|A|$ and m in the case with the 5-brane effect for $m_{3/2} = 2.0$ TeV, $\theta = \pi/2$, $\phi = \pi/4$ and $K_{5,Z\bar{Z}} = K_{5,Z} = 1$.
- Fig. 7 5-brane effects on the electroweak symmetry breaking, the constraint $m_{\tilde{\tau}_1}^2 > 0$ and the LSP for $\theta = \pi/2$ and $K_{5,Z\bar{Z}} = K_{5,Z} = 1$. Figs. 7a and 7b correspond to $\phi = \pi/4$ and $\pi/5$.
- Fig. 8 The superparticle spectra against $2\alpha Re(T)$ and π . Fig. 8a shows $2\alpha Re(T)$ -dependence of the lightest neutralino, chargino, stau and stop masses for $m_{3/2} = 1.0$ TeV, $\theta = \pi/2, \pi/4$ and $K_{5,Z\bar{Z}} = K_{5,Z} = 1$. Similarly, Fig. 8b shows ϕ -dependence of the superparticle masses for $m_{3/2} = 1.0$ TeV, $\theta = \pi/2$, $2\alpha Re(T) = 0.5$ and $K_{5,Z\bar{Z}} = K_{5,Z} = 1$.
- Fig. 9 The soft parameters, $M_{1/2}$, $|A|$ and m when we vary $K_{5,Z\bar{Z}}$. The other parameters are fixed $m_{3/2} = 2.0$ TeV, $\theta = \pi/2$, $\phi = \pi/4$, $2\alpha Re(T) = 1.0$ and $K_{5,Z} = 1$.

Fig. 10 $K_{5,Z\bar{Z}}$ -dependence of the electroweak symmetry breaking, the constraint $m_{\tilde{\tau}_1}^2 > 0$ and the LSP for $m_{3/2} = 2.0$ TeV, $\theta = \pi/2$, $\phi = \pi/4$ and $K_{5,Z} = 1$.

Fig. 11 $K_{5,Z}$ -dependence of the electroweak symmetry breaking, the constraint $m_{\tilde{\tau}_1}^2 > 0$ and the LSP for $m_{3/2} = 2.0$ TeV, $\theta = \pi/2$, $\phi = \pi/4$ and $K_{5,Z\bar{Z}} = 1$. Fig. 11a and 11b correspond to the regions with $-1.0 \leq K_{5,Z} \leq 1.0$ and $1.0 \leq K_{5,Z} \leq 2.0$, respectively.

Fig. 12 The superparticle spectrum against $K_{5,Z}$ for $m_{3/2} = 1.0$ TeV, $\theta = \pi/2$, $\pi/4$, $2\alpha Re(T) = 0.5$ and $K_{5,Z\bar{Z}} = 1$.

Fig.1




Fig.2a




Fig.2b




Fig.2c




Fig.2d




Fig.3a




Fig.3b




Fig.4a




Fig.4b




Fig.4c




Fig.5a




Fig.5b




Fig.6




Fig.7a




Fig.7b




Fig.8a




Fig.8b




Fig.9




Fig.10




Fig.11a




Fig.11b




Fig.12

